

Geographically structured genetic variation in the *Medicago lupulina*–*Ensifer* mutualism

Tia L. Harrison,^{1,2} Corlett W. Wood,¹ Katy D. Heath,³ and John R. Stinchcombe^{1,4} 

¹Department of Ecology and Evolutionary Biology, University of Toronto, 25 Willcocks Street, Toronto, Ontario, Canada M5S 3B2

²E-mail: tia.harrison@mail.utoronto.ca

³Department of Plant Biology, University of Illinois, 505 S. Goodwin Avenue, Urbana, Illinois 61801

⁴Centre for Genome Evolution and Function, University of Toronto, 25 Willcocks Street, Toronto, Ontario, Canada M5S 3B2

Received November 15, 2016

Accepted April 15, 2017

Gene flow between genetically differentiated populations can maintain variation in species interactions, especially when population structure is congruent between interacting species. However, large-scale empirical comparisons of the population structure of interacting species are rare, particularly in positive interspecific interactions (mutualisms). One agriculturally and ecologically important mutualism is the partnership between legume plants and rhizobia. Through characterizing and comparing the population genomic structure of the legume *Medicago lupulina* and two rhizobial species (*Ensifer medicae* and *E. meliloti*), we explored the spatial scale of population differentiation between interacting partners in their introduced range in North America. We found high proportions of *E. meliloti* in southeastern populations and high proportions of *E. medicae* in northwestern populations. *Medicago lupulina* and the *Ensifer* genus showed similar patterns of spatial genetic structure (isolation by distance). However, we detected no evidence of isolation by distance or population structure within either species of bacteria. Genome-wide nucleotide diversity within each of the two *Ensifer* species was low, suggesting limited introduction of strains, founder events, or severe bottlenecks. Our results suggest that there is potential for geographically structured coevolution between *M. lupulina* and the *Ensifer* genus, but not between *M. lupulina* and either *Ensifer* species.

KEY WORDS: Coevolution, gene flow, genetic differentiation, invasion, mutualism, population genetics.

The maintenance of variation within mutualistic interactions has been posed as a paradox because strong selection is expected to erode variation in mutualism-related traits (Charlesworth 1987, Heath and Stinchcombe 2013). One simple mechanism that could resolve this paradox is genetic differentiation between populations in mutualism traits, coupled with some gene flow between populations that introduces new variants. To evaluate this possibility, it is necessary to incorporate a geographic perspective into studies of mutualism to determine whether both interacting partners exhibit similar patterns of genetic structure on a landscape scale. Here, we use whole genome sequencing and genotyping-by-sequencing (GBS) to characterize patterns of genetic and geographic differentiation in the annual legume *Medicago lupulina* and its mutualistic rhizobial symbionts in their introduced North American range.

The potential for geographic structure to maintain variation in interspecific interactions is a core component of the geographic mosaic perspective on coevolution. A geographic mosaic describes a scenario where the structure and intensity of coevolution differs between populations, and is characterized by genetic differentiation between interacting populations at loci underlying coevolutionary traits, followed by gene flow that introduces new variants (Thompson 2005). Adaptive genetic divergence in coevolutionary traits can arise from interactions with genetically differentiated populations of a single partner species or turnover of partner assemblages across a focal species' range (Nagano et al. 2014; Newman et al. 2015). Formal theory and meta-analyses suggest that gene flow between genetically differentiated populations can facilitate local adaptation in host–parasite systems

by increasing within-population genetic variance (Gandon et al. 1996; Greischar and Koskella 2007; Hoeksema and Forde 2008; Gandon and Nuismer 2009). Although theoretical models indicate that geographic structure may similarly maintain genetic variance in mutualisms (Nuismer et al. 2000), empirical evidence in positive species interactions is scarce.

Gene flow between differentiated populations has the greatest potential to maintain variation in interspecific interactions when the scale of population differentiation in both partners is congruent. Although there is strong evidence of geographic variation in mutualist quality (Thrall et al., 2000, 2007), and geographic covariation in traits mediating interactions (Anderson and Johnson 2007), we lack large-scale empirical examinations of population genetic structure in interacting mutualists. The few empirical studies that have examined parallel patterns of geographic structured genetic variation in both partners report conflicting results. Anderson et al. (2004), for example, found parallel patterns of isolation by distance between carnivorous *Roridula* plants and their hemipteran mutualists, albeit at different spatial scales, and suggested that these population genetic structures could facilitate coadaptation within populations or regions. Parker and Spoerke (1998), in contrast, found no evidence of genetic structure in either the annual legume *Amphicarpea bracteata* or its nitrogen-fixing rhizobial symbionts. Béna et al. (2005) reported suggestive evidence of cospeciation between legumes in the genus *Medicago* and their rhizobial symbionts, but this genus-level analysis was not able to link phylogenetic patterns to coevolutionary processes that might have generated them.

In this study, we characterized and compared the geographic scale of genetic differentiation between the annual legume, *Medicago lupulina*, and its mutualistic nitrogen (N)-fixing bacteria, *Ensifer meliloti* and *E. medicae*, to determine whether gene flow between differentiated populations could maintain variation in this mutualism. Within the mutualism, legumes provide carbon-based rewards and shelter for the bacteria (rhizobia), whereas bacteria fix atmospheric nitrogen (N) into plant-available forms. The *Medicago–Ensifer* mutualism is characterized by considerable coevolutionary genetic variation (Heath 2010; Heath et al. 2012), and several aspects of its biology suggest that there is substantial potential for geographic structure in both partners. *Medicago lupulina* is primarily a selfer, which reduces gene flow via pollen and promotes genetic differentiation. In addition, *M. lupulina* and *Ensifer* were introduced to North America relatively recently (approximately 300 years ago) and potentially multiple times (Turkington and Cavers 1979). Multiple and separate introductions of *M. lupulina* and *Ensifer* to North America could have created the necessary geographic structure to maintain mutualism variation in its introduced range.

One challenge in evaluating the potential for geographic structure to maintain genetic variation in mutualistic traits is that

geographic structure might only be detected at specific genes involved in the mutualism. Although genetic structure at genes involved in adaptation to other aspects of the environment will contribute to population divergence, these differences will not result in divergence in mutualism-related traits or genes, except in the case of linkage disequilibrium or pleiotropy. Therefore, a rigorous test of geographic structure in mutualisms would ideally quantify patterns of structure at symbiosis genes in addition to the whole genome. The mutualism between legumes and nitrogen-fixing rhizobia is especially promising in this regard. Genes mediating the interactions have been mapped (Wernegreen and Riley 1999; Barnett et al. 2001; Markmann and Parniske 2009; Reeve et al. 2010; Oldroyd 2013; Stanton-Geddes et al. 2013; Bravo et al. 2016; Klinger et al. 2016) and it is feasible to sequence entire bacterial genomes with next-generation sequencing rather than just a handful of markers. Using both whole genome sequences and sequences of symbiotic loci such as nitrogen fixation and nodulation genes previously shown to be involved in the symbiosis between *M. lupulina* and *Ensifer* (Wernegreen and Riley, 1999; Kimbrel et al., 2013; Kawaharada et al. 2015), we looked for signals of coevolution between legumes and their rhizobia genome wide and at individual symbiotic genes.

We asked three questions about the *M. lupulina* and *Ensifer* mutualism. First, is there geographic structure in the distribution of *E. meliloti* and *E. medicae* that could facilitate differentiation of *M. lupulina* populations? Second, do symbiotic genes in rhizobia indicate alternative patterns of coevolution compared to the whole genome? Finally, is population genetic structure in *M. lupulina* aligned with *Ensifer* genetic structure such that it could promote local- or regional-scale coevolution?

Methods

STUDY SYSTEM

Medicago lupulina is a clover native to eastern Europe and western Asia and was introduced (potentially multiple times) to North America in the 1700s (Turkington and Cavers 1979). Today, *M. lupulina* is found across North America in temperate and subtropical areas, including all 50 states and most Canadian provinces (Turkington and Cavers 1979). It is primarily self-fertilizing and disperses seeds passively (Turkington and Cavers 1979; Yan et al. 2009) and consistent with this, previous studies in the native range (Europe and Asia) have found significant isolation by distance (Yan et al. 2009). *Medicago lupulina* is largely considered a weed, although it has been used as an inefficient fodder plant and was potentially introduced to North America along with agricultural crops.

Two species of *Ensifer*, free-living soil bacteria native to Europe and Asia, inhabit root nodules of *M. lupulina*: *E. meliloti* and *E. medicae*. Both can also associate with other *Medicago* species

(Prévost and Bromfield 2003). It is assumed the *Ensifer* species arrived in North America with a *Medicago* species (Turkington and Cavers 1979). *Ensifer* species associate with plants at the start of a growing season, and at nodule senescence they dissociate from the plant, dispersing into soil, where they can be redistributed due to soil disturbance and water flow (i.e., no vertical transmission). Their genomes consist of a circular chromosome (3.65 Mb) and two plasmids (~1.3 and ~1.6 Mb; Galibert et al. 2001; Reeve et al. 2010). Recombination is restricted to *Ensifer* plasmids and horizontal gene transfer can occur between plasmids of different species of *Ensifer* (Bailly et al. 2006; Epstein et al. 2012; 2014). Many genes known to be involved in the mutualism, including *nif* and *nod* genes, are found on the plasmids of *E. meliloti* and *E. medicae* (Bailly et al. 2006; 2007), whereas housekeeping genes for general bacterial functions predominate on the chromosome. Past studies have failed to detect significant genetic differentiation in *E. meliloti* and *E. medicae* populations in Mexico, suggesting high levels of gene flow in *Ensifer* populations (Silva et al. 2007).

FIELD SAMPLING

We sampled *M. lupulina* individuals opportunistically from 39 populations across a wide geographic range in southern Ontario and the northeastern United States, a subset of *M. lupulina*'s introduced range (Table S1). We randomly collected two to 10 plant individuals (spaced approximately 0.5–2 m apart) in late stages of their life cycle for both seeds and nodules. Seeds were collected in envelopes in the field and nodules were kept on the roots and placed in plastic bags at 4°C until processed. We obtained samples from 28 populations in southwestern Ontario (10–300 km apart). To study large-scale geographic patterns, we sampled an additional 11 populations along a NW to SE transect from southern Ontario to Delaware, USA, separated by up to 820 km.

MOLECULAR PROTOCOLS

We extracted rhizobia samples from one field-collected nodule per plant, and used field-collected seeds to grow plant material for DNA extraction. Full details on plant growth conditions, bacterial plating and isolation procedures, and DNA extractions can be found in the Supporting Information Appendix. In brief, we isolated one bacterial strain per plant for whole genome sequencing using the MoBio UltraClean Microbial DNA Isolation Kit (MO BIO Laboratories Inc, Carlsbad, CA), and for the plants we isolated DNA from one individual per maternal line for GBS according to the instructions of the Qiagen DNeasy Plant Tissue Mini Protocol (QIAGEN, Toronto, ON). GBS is a high-throughput and cost-efficient method of sequencing large numbers of samples. GBS is similar to restriction site-associated sequencing (RAD-seq), and uses restriction enzymes to identify single nucleotide polymorphisms across the entire genome without sequencing the

whole genome (Elshire et al. 2011). The GBS protocol is optimized for many different plant species, including *Medicago*.

We submitted 89 bacterial DNA samples to the Hospital for Sick Children (Toronto, ON, Canada) for library preparation and whole-genome sequencing on a HiSeq Illumina platform, using one lane and 2 × 100 bp reads. For *Medicago*, we submitted 190 DNA samples to Cornell University (Ithaca, NY) for GBS. The 190 DNA samples were distributed across two 96-well plates with 95 samples and one blank in each plate for the 96 multiplex GBS protocol. Cornell University prepared genomic libraries (Elshire et al. 2011) using a single digestion with EcoT22I (sequence ATGCAT). Samples were sequenced in two Illumina flow cells lanes.

BIOINFORMATICS AND SNP DISCOVERY

We aligned forward and reverse rhizobia reads to the reference genome of *E. meliloti* strain 1021 (Galibert et al. 2001; NCBI references chromosome AIL591688, plasmid a AE006469, plasmid b AL591985) and the *E. medicae* strain WSM419 (Reeve et al. 2010; NCBI references chromosome 150026743 plasmid b 150030273, plasmid a 150031715, accessory plasmid 150032810) using BWA (Li and Durbin 2009) and Stampy (Lunter and Goodson 2011) with default parameters and the bamkeep-goodreads parameter. We assigned bacterial species using a combination of the percentage of reads mapping to one reference genome, and sequences at the 16S rDNA locus (NCBI gene references 1234653 and 5324158, respectively), which differs between *E. medicae* and *E. meliloti* (Rome et al. 1997). We used Integrative Genomics Viewer (Broad Institute, Cambridge, MA) to visualize and check alignment quality (Robinson et al. 2011). In general, 69.99–94.02% (median 84.71%) of reads per sample mapped to the *E. meliloti* reference genome, and 69.32–92.48% (median 83.49%) mapped to the *E. medicae* genome.

In addition to creating a separate SNP file for each *Ensifer* species, we also created a single SNP file containing both *E. meliloti* and *E. medicae* (hereafter referred to as the "*Ensifer* genus dataset") to assess divergence between the two rhizobia. To create this file, we aligned all strains from both species to the *E. meliloti* reference genome and performed the same SNP discovery methods as performed on the *E. meliloti* species alignments (detailed below). We found shared polymorphisms between the two species and the two species were correctly identified in Structure (Fig. S1) and in Phylip (neighbor joining; Fig. 2) using this dataset (Felsenstein, 1989, Pritchard et al. 2000). To determine whether the reference genome we used influenced our results, we also aligned all the strains to the *E. medicae* reference genome. This analysis produced similar qualitative results (it correctly identified the two *Ensifer* species in Structure; Fig. 3), so we used the *E. meliloti* alignments for the combined species SNP file for the rest of our analyses.

In *Ensifer*, we used PICARD tools to format, sort, and remove duplicates in sequence alignments. We applied GATK (Broad Institute, Cambridge, MA) version 3 indel realignment and GATK Unified Genotyper SNP discovery on all bacteria alignments (McKenna et al. 2010) with ploidy set to haploid. We used the Select Variants parameter in GATK to select SNP variants only. We used standard hard filtering parameters and variant quality score recalibration on SNP discovery according to GATK Best Practices (DePristo et al. 2011; Van der Auwera et al. 2013). We filtered rhizobia SNPs for a minimum read depth (DP) of 20, a maximum DP of 226 for *E. meliloti* (230 for *E. medicae*), and a genotype quality (GP) of 30 using vcfTools (Danecek et al. 2011). We removed indels and sites with more than 10% of missing data from both *E. meliloti* and *E. medicae* data files. We identified synonymous SNPs using SnpEff (Cingolani et al. 2012a) and SnpSift (Cingolani et al. 2012b), using reference files GCA_000017145.1.22 and GCA_000006965.1.22 (for *E. medicae* and *E. meliloti*, respectively) in the prebuilt database. We used the ANN annotation parameter in SnpSift to identify SNPs as synonymous variants and missense variants.

We called *Medicago* SNPs in GBS samples by following the three-stage pipeline in the program Stacks (Catchen et al. 2011; 2013): cleaning raw data, building loci, and identifying SNPs. We trimmed reads to 64 bp and filtered reads by a phred score of 33, the default value for GSB reads sequenced on Illumina 2000/2500 machine. We built loci for *M. lupulina* using the *de novo* approach in Stacks (denovo_map command), setting the $-m$ parameter at 5, the $-M$ parameter at 1, and the $-n$ parameter at 1. In the final stage of the pipeline, we identified SNPs under the populations command by setting the $-m$ parameter at 5. We filtered SNPs by removing indels, removing sites with more than 10% of missing data, and removing sites that were less than 64 bp apart with vcfTools (Danecek et al. 2011). We also excluded nine SNPs with heterozygosity that was higher than expected under Hardy–Weinberg.

ANALYSIS OF *M. lupulina* AND *Ensifer* GENETIC STRUCTURE

We tested whether genetic distance was correlated with geographic distance (isolation by distance) in *Medicago* and *Ensifer* using Mantel tests, implemented in R (R Core Team 2016) with the ade4 package (Dray and Dufour 2007) using 100,000 randomizations. We estimated pairwise genetic distances between populations in *M. lupulina* and between individual samples in *Ensifer* because we sampled relatively few rhizobia from each population (one to three samples). For *M. lupulina*, we used SNPs to calculate pairwise F_{ST} between populations in the program Genodive (Meirmans and van Tienderen 2004) using the population F_{ST} function and 1000 permutations, including only populations that had at least two individuals in F_{ST} estimates. We converted F_{ST}

values to genetic distance values using $F_{ST}/(1 - F_{ST})$; Rousset 1997). In addition to calculating genetic distance between plant populations, we also used F -statistics to test for genetic differentiation between individuals hosting different species of bacteria, and to estimate population-level selfing rates [$s = 2F_{IS}/(1 + F_{IS})$] (Hartl and Clarke 1989). For *Ensifer*, we calculated Rousset's genetic distance between strains in the program Genepop using the combined *E. medicae* and *E. meliloti* SNP dataset (Rousset 2008). To test for isolation by distance within *Ensifer* species, we repeated this procedure separately for *E. medicae* and *E. meliloti* datasets, and also computed separate tests of isolation by distance for the chromosome and plasmid to assess structure at different components of the *Ensifer* genome.

Second, we tested for spatial genetic autocorrelation of allele frequencies in *M. lupulina*, in the *Ensifer* genus, and separately in each *Ensifer* species using GenAlEx version 6.5 (Peakall and Smouse 2006; 2012). This analysis tests against the null hypothesis that genotypes are randomly distributed in space. We binned individuals into eight distance classes of 100 km for the *M. lupulina* and *Ensifer* genus analyses, and into four distance classes of 200 km for the separate analyses of each *Ensifer* species, because our sample sizes were smaller for the latter two analyses. We tested for significant spatial autocorrelation by permuting individuals among geographic locations ($N_{\text{permutations}} = 999$) and placed confidence limits on our estimates of spatial autocorrelation using 1000 bootstrap replicates.

Finally, we tested for a geographic pattern in the distribution of the two *Ensifer* species. Because our sampling transect ran from northwest to southeast, we created a single variable representing increasing longitude and decreasing latitude by extracting the first principal component (PC1) from the latitude and longitude coordinates of our collection sites. The PC1 axis captured 90.79% of the variance in geographic location among our collection sites. We regressed the proportion of *E. meliloti* samples in a site on PC1 to identify the relationship between *Ensifer* species proportion and geographic location (R Core Team 2016). To assess whether spatial autocorrelation of plant samples impacted the results of this analysis, we randomly removed 17 Ontario populations and reran our analysis on the remaining 11 Ontario populations and the 11 American populations. We repeated this procedure 100 times, and obtained qualitatively similar results to the full dataset in all cases ($P \leq 0.0001$ in all cases), indicating that the geographic pattern in the distribution of the bacteria species is robust to our uneven geographic sampling.

ANALYSIS OF RHIZOBIAL NUCLEOTIDE DIVERSITY AND SYMBIOSIS GENES

We next looked for genetic variation between strains within the same *Ensifer* species. Specifically, we assessed nucleotide diversity within *Ensifer* species by calculating the average pairwise

nucleotide differences (π) between rhizobial samples. We extracted average pairwise nucleotide differences from *Ensifer* vcf files using a custom Python script (Python Core Team 2015). We averaged all pairwise nucleotide differences across strains to obtain π , and divided it by the number of loci (variant and non-variant) called by GATK to obtain per site values. We calculated π for the range-wide sample, and repeated this calculation including only individuals collected from southern Ontario, which are in close proximity and more likely to experience similar environmental (and potentially selective) conditions. We calculated π separately for the *Ensifer* chromosome and two plasmids and for synonymous and nonsynonymous SNPs in both species of *Ensifer*.

In addition to calculating nucleotide diversity at the genome-wide scale, we also calculated nucleotide diversity for individual genes known to be involved in the symbiosis between *M. lupulina* and *Ensifer* species (Wernegreen and Riley 1999): nodulation genes *nodA*, *nodB*, and *nodC*; and nitrogen fixation genes *nifA*, *nifB*, *nifD*, *nifE*, *nifH*, *nifK*, *nifN*, and *nifX* (NCBI gene reference numbers given in Table S2). Previous research has also identified pathogen type III effector genes as important genes in host infection (Kimbrel et al. 2013), so we calculated nucleotide diversity for two type III effector loci in *E. medicae* (Reeve et al. 2010). In addition, there is evidence that bacterial exopolysaccharides are involved in nodule formation and rhizobia infection (Kawaharada et al. 2015). We estimated nucleotide diversity in one gene (exoU glucosyltransferase) that produces exopolysaccharides in *E. meliloti* (Finan et al. 2001).

To further characterize diversity among rhizobia samples and more specifically assess how rare polymorphisms are in the rhizobia samples, we also constructed minor allele frequency spectra of the *E. medicae* of *E. meliloti* data. We removed 100% of missing data from the *E. medicae* and *E. meliloti* vcf files before calculating allele frequencies for synonymous and nonsynonymous SNPs using vcfTools (Danecek et al. 2011). We extracted the least frequent alleles from the *Ensifer* vcf files and constructed histograms of *E. medicae* and *E. meliloti* minor allele frequencies in R using the plotrix package.

COMPARISON OF *M. lupulina* AND *Ensifer* GENETIC STRUCTURE

To determine whether *M. lupulina* and *Ensifer* exhibited similar patterns of isolation by distance, we tested whether pairwise genetic distances between *M. lupulina* individuals were correlated with pairwise genetic distances between their rhizobia, using a Mantel test with 100,000 randomizations. We used *Ensifer* genus dataset (combined *E. meliloti* and *E. medicae*) to estimate individual genetic distance in *Ensifer*.

We estimated population structure among samples in *M. lupulina* and in the *Ensifer* genus using a combination of In-

Struct (Gao et al. 2007) and Structure (Pritchard et al. 2000). For *M. lupulina*, we tested for a maximum population value (K) of 5 under the admixture and population selfing rate model ($\nu = 2$) in the program InStruct (which allows for population assignments in selfing organisms). We ran two chains for each K -value with 500,000,000 repetitions and a burn-in of 200,000,000 and included no prior information. All other InStruct parameters were kept at default values. The Gelman–Rudin statistic confirmed that convergence among chains was achieved. We used the deviance information criteria (DIC) to select the value of K that provided the best fit to the data. We postprocessed Structure runs using CLUMPP (Jakobsson and Rosenberg 2007) and made plots using Distruct (Rosenberg 2004).

Before we estimated population structure in rhizobia strains using Structure, we first estimated recombination among the samples. The Structure model assumes that loci are not in linkage disequilibrium within populations (Pritchard et al. 2000), which is likely to be untrue for nonrecombining regions such as the *Ensifer* chromosome (Bailey et al. 2006). We used the program ClonalFrame (Didelot and Falush 2007) to estimate ρ/θ (number of recombination events/number of mutation events). We used VCFx software (Castelli et al. 2015) to convert our *Ensifer* genus vcf file of combined *E. meliloti* and *E. medicae* SNPs to an aligned fasta file—the input format for ClonalFrame. We performed two runs of ClonalFrame with 100,000 iterations and removed 50,000 as the burn-in. We checked for convergence using Gelman and Rubin's statistic. ClonalFrame identified a sufficiently high rate of recombination ($\rho/\theta = 1.0021$) among *Ensifer* samples to justify Structure analysis. In Structure (Pritchard et al. 2000), we performed five runs with 200,000 iterations and discarded 100,000 for the burn-in. We tested for a maximum K of 5 under a model of admixture and correlated allele frequencies. We used StrAuto to automate Structure processing of samples (Chhatre and Emerson 2017). All summary statistics (alpha, F_{ST} , and likelihood) stabilized before the end of the burn-in. We then used Structure Harvester to detect the inferred K in the likelihood data generated by the Structure tests (Earl 2012), using the ΔK approach (Evanno et al. 2005). Structure runs were postprocessed and plotted as described above.

To assess phylogenetic congruence between *Medicago* and *Ensifer*, we estimated phylogenetic relationships among individuals for the plant and the rhizobia by constructing maximum likelihood trees in RAxML (Stamatakis 2014). We used the GTRGAMMA function with 100 bootstraps to build our trees. Because we used SNP alignment files without invariable sites included, we used the ASC_string to apply an ascertainment bias correction to our dataset. We built a maximum likelihood tree for *M. lupulina* samples and the *Ensifer* genus (based on the combined *E. medicae* and *E. meliloti* SNP data). We then used the cophyloplot function and the dist.topo function in phangorn (Schliep




Figure 1. Relationship between geographic distance and genetic distance in *Medicago lupulina* populations. Each point represents a pairwise population comparison. One population was removed from the *M. lupulina* dataset because it produced an abnormally high genetic distance value when compared pairwise with other populations (population 11).

2011) in R to visualize the two trees and calculate topological distance between the trees. We also estimated separate neighbor joining trees for the *Ensifer* chromosome and two plasmids using the ape package (Paradis et al. 2004) in R to compare structure at different components of the *Ensifer* genome.

Results

M. lupulina GENETIC STRUCTURE

The *M. lupulina* sample of 190 individuals comprised 39 populations and 2349 SNPs, and exhibited a significant signal of isolation by distance (Fig. 1). The positive relationship between geographic distance and genetic distance indicates that populations farther apart are more genetically different than populations located close together. Population-level selfing rates (Table S3) were quite high on average ($s = 0.813$), which may contribute to isolation by distance in *M. lupulina*. F_{ST} between *M. lupulina* individuals hosting *E. medicae* and individuals hosting *E. meliloti* was low (0.0190 ± 0.0001) but significant ($P = 0.0010$).

There was significant spatial autocorrelation of allele frequencies in *M. lupulina* (Table S4; Fig. S4A). We found a positive spatial autocorrelation between individuals located within approximately 200 km of each other ($r \geq 0.04$, $P = 0.001$), indicating that geographically proximate individuals are more closely related than the null expectation. We found a negative spatial autocorrelation between individuals located farther than 300 km from each other ($r \leq -0.01$, $P = 0.001$), indicating that geographically distant individuals are less closely related than the

null expectation. These results are consistent with the pattern of isolation-by-distance reported above.

Ensifer GENETIC STRUCTURE

We assigned 50 rhizobia samples to *E. meliloti* and 39 samples to *E. medicae*; summary statistics on sequencing can be found in Tables S5 and S6. The 39 *E. medicae* samples were distributed among 24 populations. In this dataset, we discovered 1081 SNPs, of which 678 were synonymous and 209 nonsynonymous. The 50 *E. meliloti* sample were distributed among 28 populations, but contained approximately half the number of SNPs that *E. medicae* did (554: 234 synonymous and 176 nonsynonymous). Our *Ensifer* genus dataset (combining both *E. meliloti* and *E. medicae*) contained a total of 89 samples and 476 SNPs; this dataset contained fewer SNPs than either the *E. medicae* or *E. meliloti* datasets because it only includes sites that were genotyped in both species.

Population composition of bacteria species changed significantly with longitude and latitude. When we regressed the proportion of plants associated with *E. meliloti* on PC1, which represented increasing longitude and decreasing latitude of our sampling locations, we found a positive significant relationship ($F_{1,37} = 15.804$, $P < 0.001$). Populations in the southeast contained higher proportions of *E. meliloti*, whereas populations in the northwest contained higher proportions of *E. medicae* (Fig. 2).

We found a significant signal of isolation by distance in our *Ensifer* genus dataset (Fig. 3a), as expected given the geographic cline in their frequencies (Fig. 2). We failed to detect isolation




Figure 2. Population composition of *Ensifer meliloti* and *E. medicae* in *Medicago lupulina* populations in North America. Radius of circle corresponds to the number of *M. lupulina* samples collected in the population. Pie charts represent the proportion of plants from each population that were hosting *E. meliloti* (red), *E. medicae* (blue), and an unidentified rhizobia species (gray). Populations are numbered from south to north.




Figure 3. Relationship between geographic distance and Rousset's individual genetic distance in (a) total *Ensifer* genus dataset (*E. meliloti* and *E. medicae*), (b) *E. meliloti*, and (c) *E. medicae*. Each point represents a pairwise individual comparison.

by distance within either *Ensifer* species using whole-genome data (Fig. 3b and c). There was also no significant isolation by distance when we performed this analysis using only SNPs from the bacterial chromosome and plasmids in either *Ensifer* species (*E. medicae*: $0.23 < P < 0.65$; *E. meliloti*: $0.9 < P < 0.96$).

There was significant spatial autocorrelation in allele frequencies in the *Ensifer* genus (Table S4; Fig. S4B). We found a positive spatial autocorrelation between individuals located within approximately 300 km of each other ($r \geq 0.02$, $P \leq 0.015$), and a negative spatial autocorrelation between individuals located at

least 600 km from each other ($r \leq -0.05$, $P \leq 0.004$). These results are consistent with the pattern of isolation-by-distance reported above, in which geographically proximate individuals are more genetically similar (in this case, of the same species) and geographically distant individuals are more genetically dissimilar (i.e., of alternate bacterial species) than expected by chance. By contrast, there was no significant spatial autocorrelation of allele frequencies within either *Ensifer* species when the two species were analyzed separately (Table S4; Fig. S4C and D).

Ensifer NUCLEOTIDE DIVERSITY AND SYMBIOSIS GENES

Genome-wide nucleotide diversity values were extremely low within both *Ensifer* species in our full range dataset and reduced dataset in Ontario (Table 1). Symbiosis genes were particularly conserved (Table 2). We discovered only one to two SNPs in the *nodC* nodulation gene in both species of *Ensifer*. *NodA* and *nodB* genes contained no SNPs in either *E. medicae* or *E. meliloti*. In addition, *nifH* was the only nitrogen fixation gene that contained SNPs in both *E. medicae* and *E. meliloti*; *nifE* in *E. medicae* was the only other nitrogen fixation gene with a nucleotide diversity value greater than zero. We detected no SNPs in *E. medicae* type III effector genes or exopolysaccharide genes in *E. meliloti*, which are known to be involved in nodule formation and rhizobia infection (Kawaharada et al. 2015).

Minor allele frequency spectra showed that most minor alleles were very low in frequency in *E. meliloti* and *E. medicae* (Fig. S5). Minor alleles are all quite rare in *E. medicae* as almost all

Table 1. Nucleotide diversity (mean π) of *Ensifer medicae* and *E. meliloti* for different structures of the *Ensifer* genome.

	π Synonymous	π Nonsynonymous
Full range sample		
<i>E. medicae</i>		
Chromosome	0.0006108	0.0001117
pSMED01	0.0010950	0.0002371
pSMED02	0.0025284	0.0010754
<i>E. meliloti</i>		
Chromosome	0.0001349	0.0000312
pSymA	0.0005108	0.0003873
pSymB	0.0001449	0.0000362
Southern Ontario sample		
<i>E. medicae</i>		
Chromosome	0.0004844	0.0000931
pSMED01	0.0021592	0.0008977
pSMED02	0.0009104	0.0002091
<i>E. meliloti</i>		
Chromosome	0.0001324	0.0000283
pSymA	0.0005056	0.0003586
pSymB	0.0001338	0.0000383

Table 2. Nucleotide diversity (mean π) on nodulation genes and nitrogen fixation genes located on *Ensifer medicae* and *E. meliloti* plasmids.

	Number of SNPs	Π
<i>E. medicae</i>		
<i>nodA</i>	0	0
<i>nodB</i>	0	0
<i>nodC</i>	2	0.0000761
<i>nifA</i>	0	0
<i>nifB</i>	0	0
<i>nifD</i>	0	0
<i>nifE</i>	1	0.0000359
<i>nifH</i>	3	0.0004896
<i>nifK</i>	0	0
<i>nifN</i>	0	0
<i>nifX</i>	0	0
Type III effector 4319	0	0
Type III effector 1279	0	0
<i>E. meliloti</i>		
<i>nodA</i>	0	0
<i>nodB</i>	0	0
<i>nodC</i>	1	0.0002755
<i>nifA</i>	0	0
<i>nifB</i>	0	0
<i>nifD</i>	0	0
<i>nifE</i>	0	0
<i>nifH</i>	1	0.0001262
<i>nifK</i>	0	0
<i>nifN</i>	0	0
<i>nifX</i>	0	0
exoU glucosyltransferase	0	0

the alleles were below 5% in frequency. Minor allele frequencies in *E. meliloti* had more variation across the different frequency bins compared to *E. medicae*, but still most of the alleles were low in frequency (5%).

COMPARISON OF *M. lupulina* AND *Ensifer* GENETIC STRUCTURE

We found a significant positive relationship between *M. lupulina* genetic distance and *Ensifer* genetic distance (Fig. 4). The positive relationship indicates that as genetic divergence between plant populations increased, so did genetic differentiation between their associated rhizobia.

We compared population assignments in *Ensifer* samples to population assignments in their specific *M. lupulina* individual hosts. We identified two genetic clusters within *M. lupulina* using Instruct (Fig. 5a), using the DIC to determine which value of K provided the best fit to the data. There is a weak geographic trend of northern *M. lupulina* individuals associated with the purple




Figure 4. Relationship between *Medicago lupulina* individual genetic distance and *Ensifer* individual genetic distance. Each point represents a pairwise comparison between the genetic distance between two *M. lupulina* individuals and the genetic distance between their two corresponding rhizobia strains.




Figure 5. Population structure of (a) *Medicago lupulina* and (b) *Ensifer* genus. Black lines represent population divisions in the sample. Geographic population numbers are listed on the x-axis and are ordered from south populations to north populations.

cluster, and southern *M. lupulina* individuals associated with the yellow cluster. Similarly, Structure Harvester identified two clusters within the *Ensifer* genus dataset, corresponding to *E. medicae* and *E. meliloti* (Fig. 5b). All *E. meliloti* samples were assigned to the red population and all *E. medicae* samples were assigned to the blue population.

The maximum likelihood trees of *M. lupulina* and *Ensifer* show extensive mismatching between tree tips (Fig. 6). Plants hosting *E. medicae* and plants hosting *E. meliloti* did not group together on the *M. lupulina* tree. In addition, topological distance (the number of partitions that differ between the two trees) was high (topological distance = 140, total partitions = 140, percent differences in bipartitions between

trees = 100%). It is important to note that both trees had low bootstrap support at internal nodes. The *Ensifer* tree had particularly low bootstrap at nodes within *Ensifer* species (which could be a result of the low genetic diversity within *Ensifer* species). Therefore, mismatches between *M. lupulina* and *Ensifer* at the tree tips are likely due in part to error associated with clade assignments.

Groupings in the maximum likelihood tree of *M. lupulina* samples did not necessarily corresponded to groupings of geographic populations. The tree topology also showed large genetic distance between individuals. The tree topology for the *Ensifer* genus showed *E. medicae* and *E. meliloti* clearly separated into two groups (Fig. 6). Groupings of *Ensifer* samples in the tree did not necessarily associate with geographic location, even when




Figure 6. Phylogenetic analysis of *Medicago lupulina* (left) and *Ensifer* (right) estimated using genome-wide SNPs. Maximum likelihood trees with posterior support given at each node. Circles at nodes indicate varying bootstrap support with the colors white (< 75%), gray (> 75 < 90%), and black (> 90%). Scale bar represents the genetic distance between individuals. Number codes represent populations and individuals within populations. Individuals are also labeled for which rhizobia species they were associated with in the sample (left tree) or which rhizobia species (right tree) they were identified as (red = *Ensifer meliloti* and blue = *E. medicae*).

we constructed separate trees for the *Ensifer* chromosome and two plasmids. The chromosome and plasmid trees differed appreciably (Figs. S6 and S7). In general, the *Ensifer* tree had lower genetic distance between individuals when compared to the *M. lupulina* tree.

Discussion

Our primary goal was to characterize and compare the spatial scale of genetic differentiation in the *M. lupulina* and *Ensifer* mutualism in a portion of its introduced range in eastern North America. The dominant picture that emerges from these analyses is that there is geographic structure in the *Ensifer* genus but very little genetic variation within each *Ensifer* species. Therefore, the geographical structure of genetic variation, and potential for coevolution in this mutualism, appears mainly to be due to *M. lupulina* interacting with different bacterial species across its range, rather than genetically variable strains within a single bacterial species. Three major results emerged from our analyses, which we discuss in turn below: (1) the geographic turnover of *Ensifer* species composition in eastern North America, (2) the overall paucity of genetic variation within both *Ensifer* species, despite an extensive collection across a wide geographic range, and (3) somewhat concordant

geographic patterns of genetic variation in *M. lupulina* and the *Ensifer* genus.

GEOGRAPHIC TURNOVER OF *Ensifer* ASSEMBLAGES AND LOW GENETIC VARIATION WITHIN *Ensifer* SPECIES

We showed that there is strong geographic structure in *Ensifer* mutualism assemblages in eastern North America. The rhizobia species *E. medicae* is more common in southern Ontario, with *E. meliloti* more common in northeastern and mid-Atlantic regions in the United States. Our results corroborate previous work, which found that *E. medicae* is more abundant in southern Ontario than other *Ensifer* species (Prévost and Bromfield 2003). Surprisingly, although we sampled across a wide geographic range, there was no evidence of population structure within each *Ensifer* species. When we assessed isolation by distance separately in *E. medicae* and *E. meliloti*, we failed to detect spatial genetic structure within either rhizobia species in the chromosome or plasmids.

A previous study, which also failed to detect population genetic structure within *Ensifer* species on a large geographic scale, attributed their result to high gene flow among *Ensifer* populations (Silva et al. 2007). High gene flow may explain the lack of

genetic structure within *Ensifer* species that we observed as well. The absence of structure across large geographic distances in both studies suggests that dispersal over distances of tens or hundreds of kilometers may frequently occur in *Ensifer*. In addition to this possibility, our data suggest that an absence of genetic structure within *Ensifer* species may be due to limited genetic variation within each species. Nucleotide diversity within each species was at least one order of magnitude lower in its introduced range in North America than in its native range (Epstein et al. 2012). Moreover, we found a near total lack of variation at symbiosis loci within *Ensifer* species, indicating that the absence of genetic structure within each *Ensifer* species does not obscure a significant signal of population differentiation at mutualism-associated loci.

A combination of founder effects, genetic bottlenecks, or recent and limited introduction of bacterial strains likely explains the lack of variation within *Ensifer* species in North America. First, the *Ensifer* samples we collected could be clones of a single strain present in North America. Perhaps a single strain of each *Ensifer* species established in North America when *Ensifer* was introduced in the 1700s (Turkington and Cavers 1979). Alternatively, the strains we sampled could be recent immigrants from *Ensifer*'s native range that have recently displaced older strains. Third, the facultative nature of the *Ensifer*–*Medicago* interaction may lead to periodic bottlenecks due to strong overwinter selection in the soil that leaves behind limited strains that are capable of associating with plants the following spring. Finally, because we sampled nodules, we only sequenced rhizobium strains that are compatible with *M. lupulina*. Knowing whether the host-compatible rhizobia are only a subset of the diversity of the entire community, as in *Bradyrhizobium* (Sachs et al. 2009), would require a much larger sample of soil diversity. Nevertheless, such a pattern would simply shift the question to why there is such little nucleotide variation among just the compatible strains.

Variation in performance among partner genotypes is important for driving the evolution of partner choice, host sanctions, and cheating in mutualisms, an area that has been explored extensively in the legume–rhizobia symbiosis (Sachs and Simms 2008; Frederickson 2013; Simonsen and Stinchcombe 2014b; Jones et al. 2015). Much of the legume–rhizobia literature assumes that legume plants have a plethora of genetically distinct rhizobia strains to choose from, and that bacterial variation is abundant due to their generation time, numerical abundance, and the number of progeny produced. The relative lack of nucleotide variation within *Ensifer* species—either genome-wide, or in genes implicated in the symbiosis pathway—suggests that in North America the only genetic variation available for plants to select upon is between the two *Ensifer* species. It is possible that recent host-mediated selection reduced diversity within bacteria species, but it is unlikely that such selection would be strong enough to eliminate 99.8% of sequence variation (π values suggest a maximum of 0.1–0.2% se-

quence variation; Table 1) across a geographic range of ~800 km. Nucleotide variation may also be a poor proxy for the quantitative trait variation upon which selection acts. Experimental manipulation of the *Ensifer* symbionts is necessary to explore whether there are differences in the nitrogen fixation efficiency of the two species that might drive local adaptation in the plant host, and evaluate whether genetically divergent *M. lupulina* populations are adapted to different species of rhizobia.

Many classic coevolutionary geographic mosaics comprise only two interacting species (e.g., Brodie et al. 2002). However, geographic mosaics can also involve multispecies assemblages that change in composition across a focal species' range, a pattern documented repeatedly in plant–pollinator mutualisms (e.g., Nagano et al. 2014; Newman et al. 2015). In these systems, spatial variation in pollinator community composition drives corresponding geographic variation in selection on floral phenotypes. The turnover in *Ensifer* assemblages that we observed in the *Medicago*–rhizobia mutualism fits a multispecies view of geographic mosaics. Our data highlight why it is crucial that studies exploring geographic variation in species interactions accurately capture the species assemblages involved. Although most *M. lupulina* plants in Ontario are associated with a different *Ensifer* species than plants in the southeastern United States, we would have concluded that there is no variation in *M. lupulina*'s rhizobial partners if we had analyzed each *Ensifer* species independently.

CONCORDANT SPATIAL GENETIC STRUCTURE IN THE *M. lupulina* AND *Ensifer* MUTUALISM

A combination of population genetic analyses— isolation by distance, maximum likelihood trees, and population structure analysis—showed strong evidence of genetic differentiation in *M. lupulina* that is somewhat concordant with geographic turnover in *Ensifer* species. We found that *E. meliloti* and *E. medicae* generally occupy different portions of *M. lupulina*'s introduced range. The two *M. lupulina* InStruct clusters weakly correspond to the two *Ensifer* Structure clusters representing the two rhizobia species (Fig. 5), and our F_{ST} analysis showed significant genetic differentiation in plants hosting alternative bacterial species. Partially concordant patterns of spatial genetic variation between *Medicago* and the *Ensifer* genus indicate that gene flow could contribute to the maintenance of variation in this mutualism.

In interactions between two partners, gene flow between divergent populations can maintain variation in traits mediating the interaction in both species. In multispecies assemblages—like the *Ensifer* assemblages we documented here—the implications for the maintenance of variation are somewhat different. Gene flow between rhizobia populations is unlikely to introduce new genetic variants within each *Ensifer* species because there is no geographic structure and no genetic variation in either

E. medicae or *E. meliloti*. Instead, dispersal of *Ensifer* species between populations may maintain variation in rhizobial species diversity in North America. Turnover in *Ensifer* assemblages could contribute to the maintenance of variation in *M. lupulina*. Because *M. lupulina* interacts with two different rhizobia species in eastern North America, gene flow between plant populations partnered with alternate *Ensifer* species has the potential to introduce novel variation in plant mutualism traits. Although turnover in *Ensifer* community assemblages may contribute to the maintenance of variation in *M. lupulina* on a continental scale, it is unlikely to be the main source of genetic variation within populations because neighboring populations tend to have the same species of *Ensifer*.

There is suggestive evidence that genetic differentiation among *Medicago* populations may arise in part from geographically structured coevolution with *Ensifer* assemblages. Béna et al. (2005) found evidence that geographically structured diversity in rhizobia potentially influenced geographically structured diversity in the *Medicago* genus in its native Eurasian range. Population genetic differentiation in *Medicago* could result from adaptation to local strains that differ in nitrogen fixation effectiveness. The *E. medicae* laboratory strain WSM419 is a more effective mutualist than the laboratory strain *E. meliloti* 1021 (Terpolilli et al. 2008), which if generally true of these species, suggests that the *Ensifer* species common in southern Ontario populations is more effective than the *Ensifer* species common in the southeastern United States. However, it is not necessarily appropriate to extrapolate these laboratory results to genetically heterogeneous natural populations, given that Béna et al. (2005) showed that rhizobia effectiveness is contingent on the specific legume host, and Terpolilli et al. (2008) evaluated the *Ensifer* species with a single *M. truncatula* genotype.

Concordant genetic structure in interacting species may not arise from coevolutionary processes that maintain genetic variation and facilitate future coevolution. The genetic differences between *M. lupulina* populations and geographic turnover in *Ensifer* assemblages could be due to several other processes, including multiple introductions to North America, adaptation to other aspects of the environment, or neutral forces. Local adaptation to the substantial climatic differences between southern Ontario and the southeastern United States (e.g., temperature, precipitation) could contribute to geographic structure in both *Medicago* and *Ensifer*. In addition, *Ensifer* associations with other *Medicago* species in North America, such as *M. sativa* and *M. polymorpha* (Rome et al. 1996; Béna et al. 2005), could be driving large-scale patterns in *Ensifer* species distribution. Genetic structure in *M. lupulina* in its native range has been attributed to self-fertilization (Yan et al. 2009), and likely contributes to the isolation by distance we observed as well. Evaluating the mechanisms behind the geographic trends that we observed is a separate question from the maintenance of genetic variation that ultimately requires

manipulative field experiments that are logistically challenging to perform with bacteria (but see Simonsen and Stinchcombe, 2014a). Despite these alternative explanations for the somewhat concordant patterns of geographic structure in *M. lupulina* and its rhizobial mutualist *Ensifer*, the significant potential for coevolution between *M. lupulina* and *Ensifer* assemblages we discovered in this study is worth further investigation. Future work involving experiments testing local adaptation of *M. lupulina* plants to its local *Ensifer* species could reveal additional evidence of coevolution in this system in the its introduced range in North America.

Conclusions and Prospects

Comparing spatial genetic structure and genome-wide variation in mutualist partners is an effective approach to determine the potential scale of coevolution between interacting species. Given the relative lack of genome-wide variation within *E. medicae* and *E. meliloti*, differences between *Ensifer* species are the only potential source of coevolutionary selection acting on *M. lupulina*. Our study shows how comparing geographic variation in two mutualists is important to understand how variation may be maintained in mutualisms, especially in introduced ranges where processes such as gene flow, bottlenecks, and multiple introductions can complicate mutualist interactions.

AUTHOR CONTRIBUTIONS

TLH, KDH, and JRS designed the research project. TLH, CWW, KDH, and JRS wrote the manuscript. TLH and JRS collected the genomic data. TLH, CWW, KDH, and JRS analyzed and interpreted the genomic data.

ACKNOWLEDGMENTS

We thank Stephen Wright, Nicole Mideo, Benjamin Gilbert, and Megan Frederickson for comments and discussion, Andrew Hall and Bruce Petrie for plant growth assistance, and Maggie Bartkowska, Adriana Salcedo, and Billie Gould for bioinformatics advice. Our work is supported by NSERC Discovery Grants and Graduate Scholarships (JRS, TLH), an EEB Departmental Post-Doc Fellowship (CWW), and the National Science Foundation (KDH). We have no conflict of interest.

DATA ACCESSIBILITY

DNA sequences are uploaded to the Sequence Read Archive of NCBI (*M. lupulina* Bioproject PRJNA378842 and Biosample SAMN06562075; *Ensifer* Bioproject PRJNA378842 and Biosample SAMN06562698). VCF files and python scripts are uploaded to Dryad, <https://doi.org/10.5061/dryad.77k64> (doi:10.5061/dryad.77k64). Sampling locations are available in Table S1.

LITERATURE CITED

- Anderson, B., and S. D. Johnson. 2007. The geographic mosaic of coevolution in a plant-pollinator mutualism. *Evolution* 62:220–225.
- Anderson, B., I. Olivieri, M. Lourmas, and B. A. Stewart. 2004. Comparative population genetic structures and local adaptation of two mutualists. *Evolution* 58:1730–1747.

- Baillly, X., I. Olivieri, S. De Mita, J.C. Cleyet-Marel, and G. Béna. 2006. Recombination and selection shape the molecular diversity pattern of nitrogen-fixing *Sinorhizobium* sp. associated to *Medicago*. *Mol. Ecol.* 15:2719–2734.
- Baillly, X., I. Olivieri, B. Brunel, J. C. Cleyet-Marel, and G. Béna. 2007. Horizontal gene transfer and homologous recombination drive the evolution of the nitrogen-fixing symbionts of *Medicago* species. *J. Bacteriol.* 189:5223–5236.
- Barnett, M. J., R. F. Fisher, T. Jones, C. Komp, A. P. Abola, F. Barloy-Hubler, L. Bowser, D. Capela, F. Galibert, J. Gouzy, et al. 2001. Nucleotide sequence and predicted functions of the entire *Sinorhizobium meliloti* pSymA megaplasmid. *Proc. Natl. Acad. Sci. USA* 98:9883–9888.
- Béna, G., A. Lyet, T. Huguet, and I. Olivieri. 2005. *Medicago*—*Sinorhizobium* symbiotic specificity evolution and the geographic expansion of *Medicago*. *J. Evol. Biol.* 18:1547–1558.
- Bravo, A., T. York, N. Pumplun, L. A. Mueller, and M. J. Harrison. 2016. Genes conserved for arbuscular mycorrhizal symbiosis identified through phylogenomics. *Nat. Plants* 2:15208.
- Brodie, E. D. J., B. J. Ridenhour, and E. D. Brodie III. 2002. The evolutionary response of predators to dangerous prey: hotspots and coldspots in the geographic mosaic of coevolution between garter snakes and newts. *Evolution* 56:2067–2082.
- Castelli, E. C., C. T. Mendes-Junior, A. Sabbagh, I. O. P. Porto, A. Garcia, J. Ramalho, T. H. A. Lima, J. D. Massaro, F. C. Dias, C. V. A. Collares, et al. 2015. HLA-E coding and 3' untranslated region variability determined by next-generation sequencing in two West-African population samples. *Hum. Immunol.* 76:945–953.
- Catchen, J., P. A. Hohenlohe, S. Bassham, A. Amores, and W. A. Cresko. 2013. Stacks: an analysis tool set for population genomics. *Mol. Ecol.* 22:3124–3140.
- Catchen, J. M., A. Amores, P. Hohenlohe, W. Cresko, and J. H. Postlethwait. 2011. Stacks: building and genotyping Loci de novo from short-read sequences. *G3* 1:171–182.
- Charlesworth, B. 1987. The heritability of fitness. Pp. 21–40 in J. Bradbury and M. B. Anderson, eds. *Sexual selection: testing the alternatives*. John Wiley & Sons, London, U.K.
- Chhatre, V. E., and K. J. Emerson. 2017. StrAuto: automation and parallelization of STRUCTURE analysis. *BMC Bioinform.* 18:192.
- Cingolani, P., A. Platts, L. L. Wang, M. Coon, T. Nguyen, L. Wang, S. J. Land, X. Lu, and D. M. Ruden. 2012a. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-3. *Fly (Austin)* 6:80–92.
- Cingolani, P., V. M. Patel, M. Coon, T. Nguyen, S. J. Land, D. M. Ruden, and X. Lu. 2012b. Using *Drosophila melanogaster* as a model for genotoxic chemical mutational studies with a new program, SnpSift. *Front. Genet.* 3:35.
- Danecek, P., A. Auton, G. Abecasis, C. A. Albers, E. Banks, M. A. DePristo, R. E. Handsaker, G. Lunter, G. T. Marth, S. T. Sherry, et al. 2011. The variant call format and VCFtools. *Bioinformatics* 27:2156–2158.
- DePristo, M. A., E. Banks, R. Poplin, K. V. Garimella, J. R. Maguire, C. Hartl, A. A. Philippakis, G. del Angel, M. A. Rivas, M. Hanna, et al. 2011. A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nat. Genet.* 43:491–498.
- Didelot, X., and D. Falush. 2007. Inference of bacterial microevolution using multilocus sequence data. *Genetics* 175:1251–1266.
- Dray, S., and A. B. Dufour. 2007. The ade4 package: implementing the duality diagram for ecologists. *J. Stat. Softw.* 22:1–20
- Earl, D. A. 2012. STRUCTURE HARVESTER: a website and program for visualizing STRUCTURE output and implementing the Evanno method. *Conserv. Genet. Resour.* 4:359–361.
- Elshire, R. J., J. C. Glaubitz, Q. Sun, J. A. Poland, K. Kawamoto, E. S. Buckler, and S. E. Mitchell. 2011. A robust, simple genotyping-by-sequencing (GBS) approach for high diversity species. *PLoS ONE* 6:e19379.
- Epstein, B., A. Branca, J. Mudge, A. K. Bharti, R. Briskine, A. D. Farmer, M. Sugawara, N. D. Young, M. J. Sadowsky, and P. Tiffin. 2012. Population genomics of the facultatively mutualistic bacteria *Sinorhizobium meliloti* and *S. medicae*. *PLoS Genet* 8:e1002868.
- Epstein, B., M. J. Sadowsky, and P. Tiffin. 2014. Selection on horizontally transferred and duplicated genes in *sinorhizobium (ensifer)*, the root-nodule symbionts of *Medicago*. *Genome Biol. Evol.* 6:1199–1209.
- Evanno, G., S. Regnaut, and J. Goudet. 2005. Detecting the number of clusters of individuals using the software STRUCTURE: a simulation study. *Mol. Ecol.* 14:2611–2620.
- Felsenstein J. 1989. PHYLIP—phylogeny inference package (version 3.2). *Cladistics* 5:164–166.
- Finan, T. M., S. Weidner, K. Wong, J. Buhrmester, P. Chain, F. J. Vorholter, I. Hernandez-Lucas, A. Becker, A. Cowie, J. Gouzy, et al. 2001. The complete sequence of the 1,683-kb pSymB megaplasmid from the N2-fixing endosymbiont *Sinorhizobium meliloti*. *Proc. Natl. Acad. Sci. USA* 98:9889–9894.
- Frederickson, M. E. 2013. Rethinking mutualism stability: cheaters and the evolution of sanctions. *Q. Rev. Biol.* 88:269–295.
- Galibert, F., T. M. Finan, S. R. Long, A. Puhler, P. Abola, F. Ampe, F. Barloy-Hubler, M. J. Barnett, A. Becker, P. Boistard et al. 2001. The composite genome of the legume symbiont *Sinorhizobium meliloti*. *Science* 293:668–672.
- Gandon, S., and S. L. Nuismer. 2009. Interactions between genetic drift, gene flow, and selection mosaics drive parasite local adaptation. *Am. Nat.* 173:212–224.
- Gandon, S., Y. Capowiez, Y. Dubois, Y. Michalakakis, and I. Olivieri. 1996. Local adaptation and gene-for-gene coevolution in a metapopulation model. *Proc. Biol. Sci.* 263:1003–1009.
- Gao, H., S. Williamson, and C. D. Bustamante. 2007. A Markov Chain Monte Carlo approach for joint inference of population structure and inbreeding rates from multilocus genotype data. *Genetics* 176:1635–1651.
- Greischar, M. A., and B. Koskella. 2007. A synthesis of experimental work on parasite local adaptation. *Ecol. Lett.* 10:418–434.
- Hartl, D. L., and A. G. Clarke. 1989. *Principles of population genetics*. Sinauer Associates, Sunderland, MA.
- Heath, K. D. 2010. Intergenomic epistasis and coevolutionary constraint in a plant-rhizobium mutualism. *Evolution* 64:1446–1458.
- Heath, K. D., P. V. Burke, and J. R. Stinchcombe. 2012. Coevolutionary genetic variation in the legume-rhizobium transcriptome. *Mol. Ecol.* 21:4735–4747.
- Heath, K. D., and J. R. Stinchcombe. 2013. Explaining mutualism variation: a new evolutionary paradox? *Evolution* 68:309–317.
- Hoeksema, J. D., and S. E. Forde. 2008. A meta-analysis of factors affecting local adaptation between interacting species. *Am. Nat.* 171:275–290.
- Jakobsson, M., and N. A. Rosenberg. 2007. CLUMPP: a cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. *Bioinformatics* 23:1801–1806.
- Jones, E. I., M. E. Afkhami, E. Akçay, J. L. Bronstein, R. Bshary, M. E. Frederickson, K. D. Heath, J. D. Hoeksema, J. H. Ness, M. S. Pankey, et al. 2015. Cheaters must prosper: reconciling theoretical and empirical perspectives on cheating in mutualism. *Ecol. Lett.* 18:1270–1284.
- Kawaharada, Y., S. Kelly, M. W. Nielsen, C. T. Hjuler, K. Gysel, A. Muszyński, R. W. Carlson, M. B. Thygesen, N. Sandal, M. H. Asmussen, et al. 2015. Receptor-mediated exopolysaccharide perception controls bacterial infection. *Nature* 523:308–312.
- Kimbrel, J. A., W. J. Thomas, Y. Jiang, A. L. Creason, C. A. Thireault, J. L. Sachs, and J. H. Chang. 2013. Mutualistic co-evolution of type III

- effector genes in *Sinorhizobium fredii* and *Bradyrhizobium japonicum*. *PLoS Pathog.* 9:e1003204.
- Klinger, C. R., J. A. Lau, and K. D. Heath. 2016. Ecological genomics of mutualism decline in nitrogen-fixing bacteria. *Proc. Biol. Sci.* 283:20152563.
- Li, H., and R. Durbin. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25:1754–1760.
- Lunter, G., and M. Goodson. 2011. Stampy: a statistical algorithm for sensitive and fast mapping of Illumina sequence reads. *Genome Res.* 21:936–939.
- Markmann, K., and M. Parniske. 2009. Evolution of root endosymbiosis with bacteria: how novel are nodules? *Trends Plant Sci.* 14:77–86.
- McKenna, A., M. Hanna, E. Banks, A. Sivachenko, K. Cibulskis, A. Kernysky, K. Garimella, D. Altshuler, S. Gabriel, M. Daly, et al. 2010. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res.* 20:1297–1303.
- Meirmans, P. G., and P. H. van Tienderen. 2004. genotype and genodive: two programs for the analysis of genetic diversity of asexual organisms. *Mol. Ecol. Notes* 4:792–794.
- Nagano, Y., K. Abe, T. Kitazawa, M. Hattori, A. S. Hirao, and T. Itino. 2014. Changes in pollinator fauna affect altitudinal variation of floral size in a bumblebee-pollinated herb. *Ecol. Evol.* 4:3395–3407.
- Newman, E., J. Manning, and B. Anderson. 2015. Local adaptation: mechanical fit between floral ecotypes of *Nerine humilis* (*Amaryllidaceae*) and pollinator communities. *Evolution* 69:2262–2275.
- Nuismer, S. L., J. N. Thompson, and R. Gomulkiewicz. 2000. Coevolutionary clines across selection mosaics. *Evolution* 54:1102–1115.
- Oldroyd, G. E. D. 2013. Speak, friend, and enter: signalling systems that promote beneficial symbiotic associations in plants. *Nat. Rev. Micro.* 11:252–263.
- Paradis, E., J. Claude, and K. Strimmer. 2004. APE: analyses of phylogenetics and evolution in R language. *Bioinformatics* 20:289–290.
- Peakall, R., and P. E. Smouse. 2006. GENALEX 6: genetic analysis in Excel. Population genetic software for teaching and research. *Mol. Ecol. Notes* 6:288–295.
- Peakall, R., and P. E. Smouse. 2012. GenAIEx 6.5: genetic analysis in Excel. Population genetic software for teaching and research—an update. *Bioinformatics* 28:2537–2539.
- Parker, M. A., and J. M. Spoerke. 1998. Geographic structure of lineage associations in a plant-bacterial mutualism. *J. Evol. Biol.* 11:549–562.
- Prévost, D., and E. S. P. Bromfield. 2003. Diversity of symbiotic rhizobia resident in Canadian soils. *Can. J. Soil. Sci.* 83:311–319.
- Pritchard, J. K., M. Stephens, and P. Donnelly. 2000. Inference of population structure using multilocus genotype data. *Genetics* 155:945–959.
- Python Core Team. 2015. Python: A dynamic, open source programming language. Python Software Foundation. Available at <https://www.python.org/>.
- R Core Team. 2016. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available at <http://www.R-project.org>.
- Reeve, W., P. Chain, G. O'Hara, J. Ardley, K. Nandesena, L. Bräu, R. Tiwari, S. Malfatti, H. Kiss, A. Lapidus, et al. 2010. Complete genome sequence of the *Medicago* microsymbiont *Ensifer* (*Sinorhizobium*) *medicae* strain WSM419. *Stand Genomic Sci.* 2:77–86.
- Robinson, J. T., H. Thorvaldsdóttir, W. Winckler, M. Guttman, E. S. Lander, G. Getz, and J. P. Mesirov. 2011. Integrative genomics viewer. *Nat. Biotechnol.* 29:24–26.
- Rome, S., M. P. Fernandez, B. Brunel, P. Normand, and J. C. Cleyet-Marel. 1996. *Sinorhizobium medicae* sp. nov., isolated from annual *Medicago* spp. *Int. J. Syst. Bacteriol.* 46:972–980.
- Rome, S., J. C. Cleyet-Marel, L. A. Materon, P. Normand, and B. Brunel. 1997. Rapid identification of *Medicago* nodulating strains by using two oligonucleotide probes complementary to 16S rDNA sequences. *Can. J. Microbiol.* 43:854–861.
- Rosenberg, N. A. 2004. DISTRUCT: a program for the graphical display of population structure. *Mol. Ecol. Notes* 4:137–138.
- Rousset, F. 1997. Genetic differentiation and estimation of gene flow from F-statistics under isolation by distance. *Genetics* 145:1219–1228.
- . 2008. GENEPOP'007: a complete re-implementation of the genepop software for Windows and Linux. *Mol. Ecol. Resour.* 8:103–106.
- Sachs, J. L., and E. L. Simms. 2008. The origins of uncooperative rhizobia. *Oikos* 117:961–966.
- Sachs, J. L., S. W. Kembel, A. H. Lau, and E. L. Simms. 2009. In situ phylogenetic structure and diversity of wild *Bradyrhizobium* communities. *Appl. Environ. Microbiol.* 75:4727–4735.
- Schliep, K. P. 2011. phangorn: phylogenetic analysis in R. *Bioinformatics* 27:592–593.
- Silva, C., F. L. Kan, and E. Martínez-Romero. 2007. Population genetic structure of *Sinorhizobium meliloti* and *S. medicae* isolated from nodules of *Medicago* spp. in Mexico. *FEMS Microbiol. Ecol.* 60:477–489.
- Simonsen, A. K., and J. R. Stinchcombe. 2014a. Herbivory eliminates fitness costs of mutualism exploiters. *New Phytol.* 202:651–661.
- . 2014b. Standing genetic variation in host preference for mutualist microbial symbionts. *Proc. Biol. Sci.* 281:20142036.
- Stamatakis, A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30:1312–1313.
- Stanton-Geddes, J., T. Paape, B. Epstein, R. Briskine, J. Yoder, J. Mudge, A. K. Bharti, A. D. Farmer, P. Zhou, R. Denny, et al. 2013. Candidate genes and genetic architecture of symbiotic and agronomic traits revealed by whole-genome, sequence-based association genetics in *Medicago truncatula*. *PLoS ONE* 8:e65688.
- Terpolilli, J. J., G. W. O'Hara, R. P. Tiwari, M. J. Dilworth, and J. G. Howieson. 2008. The model legume *Medicago truncatula* A17 is poorly matched for N₂ fixation with the sequenced microsymbiont *Sinorhizobium meliloti* 1021. *New Phytol.* 179:62–66.
- Thompson, J. N. 2005. The geographic mosaic of coevolution. 1st ed. Univ. of Chicago Press, Chicago, IL.
- Thrall, P. H., J. J. Burdon, and M. J. Woods. 2000. Variation in the effectiveness of symbiotic associations between native rhizobia and temperate Australian legumes: interactions within and between genera. *J. Appl. Ecol.* 37:52–65.
- Thrall, P. H., M. E. Hochberg, J. J. Burdon, and J. D. Bever. 2007. Coevolution of symbiotic mutualists and parasites in a community context. *Trends Ecol. Evol.* 22:120–126.
- Turkington, R., and P. B. Cavers. 1979. The biology of Canadian weeds, *Medicago lupulina*. *Can. J. Plant Sci.* 59:99–110.
- Van der Auwera, G. A., M. O. Carneiro, C. Hartl, R. Poplin, G. del Angel, A. Levy-Moonshine, T. Jordan, K. Shakir, D. Roazen, J. Thibault, et al. 2013. From FastQ data to high confidence variant calls: the Genome Analysis Toolkit best practices pipeline. *Curr. Protoc. Bioinform.* 11:11.10.1–11.10.33.
- Wernegreen, J. J., and M. A. Riley. 1999. Comparison of the evolutionary dynamics of symbiotic and housekeeping loci: a case for the genetic coherence of rhizobial lineages. *Mol. Biol. Evol.* 1:98–113.
- Yan, J., H. J. Chu, H. C. Wang, J.-Q. Li, and T. Sang. 2009. Population genetic structure of two *Medicago* species shaped by distinct life form, mating system and seed dispersal. *Ann. Bot.* 103:825–834.

Associate Editor: M. Neiman
 Handling Editor: M. Noor

Supporting Information

Additional Supporting Information may be found in the online version of this article at the publisher's website:

Appendix. Plant Growth and Bacterial Isolation Procedures.

Table S1. Locations of *Medicago lupulina* and *Ensifer* populations used in population genetic analysis.

Table S2. NCBI gene reference numbers for nodulation and nitrogen fixation genes in *Ensifer meliloti* and *E. medicae* used in nucleotide diversity analysis.

Table S3. Heterozygosity statistics and selfing rate estimates of *Medicago lupulina* populations.

Table S4. Spatial genetic autocorrelation analyses in *Medicago lupulina*, in all *Ensifer* samples, and in each *Ensifer* species separately.

Table S5. Summary statistics on alignment of rhizobia samples to the *Ensifer meliloti* reference genome.

Table S6. Summary statistics on alignment of rhizobia samples to the *Ensifer medicae* reference genome.

Figure S1. Structure plot of all *Ensifer* samples.

Figure S2. Neighbor-joining tree of all *Ensifer* samples.

Figure S3. Structure plot of all *Ensifer* samples.

Figure S4. Spatial genetic autocorrelation analysis.

Figure S5. Minor allele frequency plots of (a) *Ensifer meliloti*, and (b) *E. medicae*.

Figure S6. Phylogenetic analysis of *Ensifer meliloti* genome structures.

Figure S7. Phylogenetic analysis of *Ensifer medicae* genome structures.